

Theory of Semi- and Nonleptonic Decays of Heavy Mesons*

Patricia Ball

Physik-Department, TU München, D-85747 Garching, Germany

September 7, 1994

Abstract:

I review some of the recent developments in the theoretical description of weak inclusive decays of heavy mesons. The topics cover the value of $|V_{cb}|$ as extracted from semileptonic inclusive decays and a short discussion of the theoretical errors. I also present the results of a recent calculation of next-to-leading order corrections to nonleptonic inclusive B decays which allows an improved prediction of the semileptonic branching ratio of B mesons.

1 Outline of Theoretical Foundations

During the recent two years, the theoretical description of inclusive decays of heavy hadrons has experienced considerable progress. For quark masses $m_Q \gg \Lambda_{\text{QCD}}$, the well-known short-distance expansion technique yields an expansion in inverse powers of the heavy quark mass, the so-called heavy quark expansion (HQE) [1]. The starting point for the HQE of, e.g., the decay rate of a B meson into a final state X is its representation as imaginary part of the relevant forward-scattering amplitude:

$$\Gamma(B \rightarrow X) = \frac{1}{m_B} \text{Im} i \int d^4x \langle B | T \mathcal{L}_W(x) \mathcal{L}_W(0) | B \rangle. \quad (1)$$

Here $\mathcal{L}_W$ is the effective weak Lagrangian mediating the decay $B \rightarrow X$. As shown in [1], for a very heavy b quark mass m_b , a short distance expansion of Eq. (1) yields:

$$\begin{aligned} \Gamma(B \rightarrow X) = & \frac{G_F^2 m_b^5}{192\pi^3} |\text{CKM}|^2 \left\{ C_0^{(X)}(\alpha_s(\mu)) \frac{\langle B | \bar{b}b | B \rangle}{2m_B} \right. \\ & \left. + C_2^{(X)}(\alpha_s(\mu)) \frac{1}{m_b^2} \frac{\langle B | \bar{b}g_s \sigma_{\mu\nu} F^{\mu\nu} b | B \rangle}{2m_B} + \mathcal{O}\left(\frac{1}{m_b^3}\right) \right\}. \end{aligned} \quad (2)$$

Here CKM denotes the appropriate CKM matrix elements, the $C_i^{(X)}$ are short-distance Wilson-coefficients, which depend on the parton model process underlying the decay $B \rightarrow X$. b is the b quark field in full QCD, m_B is the mass of the B meson and $F^{\mu\nu}$ the gluonic field-strength tensor.

*Invited talk given at ICHEP 94, Glasgow, Scotland, July 1994; to appear in the Proceedings.

Without going into too much details, let me mention just a few general features of the above expansion. First we remark that HQE strongly resembles deep inelastic scattering with the difference that the value of the expansion parameter m_b is fixed and cannot be controlled by the experimenter, so that higher twist effects in $1/m_b$ are important. On the other hand, the corresponding hadronic matrix elements can be expressed as moments of a universal distribution function [2, 3] and are thus measurable, at least in principle, cf. [4].

The first term in the above series, $\langle B|\bar{b}b|B\rangle$, just reproduces the free quark decay process; non-perturbative corrections to that picture are suppressed by terms of $\mathcal{O}(1/m_b^2)$ or higher. There are no terms of order $1/m_b$, since all possible gauge-invariant operators of suitable dimension either vanish or can be reduced to $\bar{b}b$ by the equations of motion.

The Wilson coefficients $C_j^{(X)}(\alpha_s(\mu))$ encode the short distance behaviour of Eq. (1) and are calculable within perturbation theory; they depend on the parton content of X, the renormalization scale μ and the renormalization scheme. In particular, $C_0^{(X)}$ contains the radiative corrections to the free quark decay and has been studied for various processes [5, 6, 7, 8, 9].

Let me also mention some words of caution. In order to separate clearly the expansion in $1/m_Q$ from the one in α_s , the HQE has to be done in terms of the *pole mass*. This mass definition, however, is unphysical and contradicts the confinement property of QCD. It has been shown, that this contradiction reflects itself in an intrinsic ambiguity of the definition of the pole mass which is said to be caused by a “renormalon” [10]. Although it was shown in [11] that the renormalon cancels in the semileptonic rate, at least up to order $1/m_b^2$, there is still a number of questions to be answered as far as nonleptonic and exclusive decays are concerned.

2 Applications I: $|V_{cb}|$ from Semileptonic Inclusive Decays

One immediate application of the HQE is the extraction of $|V_{cb}|$ from semileptonic inclusive decays. The decay rate $\Gamma(B \rightarrow X_c e \nu)$, of the generic form of Eq. (2), contains to order $1/m_b^2$ five unknown parameters: V_{cb} , m_b , m_c and two hadronic matrix elements:

$$\begin{aligned} 2m_B\lambda_1 &= \langle B|\bar{b}_v(iD)^2b_v|B\rangle, \\ 6m_B\lambda_2 &= \langle B|\bar{b}_v\frac{g}{2}\sigma_{\mu\nu}F^{\mu\nu}b_v|B\rangle, \end{aligned} \quad (3)$$

where b_v is defined as $b_v = e^{im_b vx}b$ and v_μ is the four-velocity of the B meson.

Whereas λ_2 is directly related to the observable spectrum of beautiful mesons,

$$\lambda_2 \approx \frac{1}{4}(m_{B^*}^2 - m_B^2) = 0.12 \text{ GeV}^2, \quad (4)$$

the quantity λ_1 is difficult to measure, cf. [4]. Physically, $-\lambda_1/(2m_b)$ is just the average kinetic energy of the b quark inside the meson. At present, only a QCD sum rule estimate is

available, according to which $\lambda_1 \simeq -0.6 \text{ GeV}^2$ [12]. This result has been met with caution (see, e.g. [13]), since it corresponds in fact to a surprisingly large momentum of the b quark inside the meson of order (700–800) MeV. However, in a recent series of papers, cf. [4, 3], an upper bound on λ_1 was derived, to wit $\lambda_1 \leq -0.4 \text{ GeV}^2$, which is in nice agreement with the QCD sum rule prediction. For a further discussion of the present status of λ_1 , I refer to [14].

The next step is to fix m_b and m_c . Here one makes use of the fact that in the framework of HQE the *difference* between m_b and m_c is given by

$$m_b - m_c = m_B - m_D + \frac{\lambda_1 + 3\lambda_2}{2} \left(\frac{1}{m_b} - \frac{1}{m_c} \right) + \mathcal{O}\left(\frac{1}{m_Q^2}\right). \quad (5)$$

Thus either m_b or m_c remain to be fixed. Shifman et al. [15] took $m_b = (4.8 \pm 0.1) \text{ GeV}$ from spectroscopy. With $\lambda_1 = -0.5 \text{ GeV}^2$ and for a B lifetime $\tau_B = 1.49 \text{ ps}$, they get $|V_{cb}| = 0.0415$. In [16, 17, 18, 19] m_c was determined from the experimental value of $B(D \rightarrow X e \nu)$ via Eq. (2) with m_b replaced by m_c . Although this procedure has the advantage that both m_b and m_c are obtained by the same method, the validity of the HQE for the charm quark is not beyond controversy. The b quark mass obtained is typically larger than 5 GeV and thus considerably larger as m_b from spectroscopy. On the other hand, the b quark mass used by Shifman et al. seems to underestimate the intrinsic renormalon ambiguity mentioned in the last section. But even using the same method, [16] and [17] obtain different results: $|V_{cb}| = 0.046 \pm 0.008$ and ≈ 0.042 , respectively (both numbers rescaled using $\tau_B = 1.49 \text{ ps}$). Ref. [19] takes into account the scheme-dependence of the free quark decay contribution and using running $\overline{\text{MS}}$ masses instead of pole masses obtains $|V_{cb}| = 0.036 \pm 0.005$. Since the decay rates only differ in terms of $\mathcal{O}(\alpha_s^2)$, these results show that at present the theoretical error due to scale- and scheme-dependence is of paramount importance in exploiting inclusive decays of heavy mesons and for $|V_{cb}|$ amounts to nearly 20%.

3 Applications II: the Semileptonic Branching Ratio of B Mesons

The semileptonic branching ratio of B mesons is defined by

$$B(B \rightarrow X e \nu) = \frac{\Gamma(B \rightarrow X e \nu)}{\Gamma_{tot}} \quad (6)$$

with

$$\Gamma_{tot} = \sum_{\ell=e, \mu, \tau} \Gamma(B \rightarrow X \ell \nu_\ell) + \Gamma(B \rightarrow X_c) + \Gamma(B \rightarrow X_{c\bar{c}}). \quad (7)$$

The explicit formulas for the decay rates can be found in [1]. The radiative corrections to $\mathcal{O}(\alpha_s)$ to the semileptonic decay $b \rightarrow c e \nu$ were calculated in [5, 7, 8], the corrections




Figure 1: $B(B \rightarrow X e \nu)$ as function of the pole mass m_b for $\mu = m_b$. Solid line: B calculated using pole masses, dashed line: B calculated using running $\overline{\text{MS}}$ quark masses.

$\alpha_s(m_Z)$	Parton Model [22]	HQE [23]	HQE [9, 24]	
	pole masses	pole masses	pole masses	$\overline{\text{MS}}$ masses
0.110	0.132	0.130	0.121	0.111
0.117	0.128	0.126	0.116	0.103
0.124	0.124	0.121	0.111	0.097

Table 1: $B(B \rightarrow X e \nu)$ in different models depending on $\alpha_s(m_Z)$. Input parameters: $m_b = 4.8 \text{ GeV}$, $m_c = 1.3 \text{ GeV}$ (pole masses) corresponding to $\lambda_1 = -0.6 \text{ GeV}^2$. Renormalization scale: $\mu = m_b$. In the phase-space factor of $\Gamma(b \rightarrow ccs)$ $m_s = 0.2 \text{ GeV}$ is used. Only in [9, 24] full radiative corrections to the nonleptonic B decay modes were taken into account.

to $b \rightarrow cud$ with a massless c quark in [6], with a massive c quark in [9]; the complete corrections to $b \rightarrow ccs$ are not known to date, but can partly be obtained from Ref. [7].

The experimental branching ratios are $B(B \rightarrow X e \nu) = (10.43 \pm 0.24)\%$ (world average) [20] and $B(B^0 \rightarrow X e \nu) = (10.9 \pm 0.7 \pm 1.1)\%$, the most recent result obtained by CLEO [21].

There exists a number of theoretical analyses of $B(B \rightarrow X e \nu)$ using different methods and obtaining different results. In [22], e.g., the branching ratio was investigated in a purely perturbative framework using the full radiative corrections for $b \rightarrow ce\nu$ [5, 8], but with $m_c = 0$ in the corrections to $b \rightarrow cud$ and $b \rightarrow ccs$ [6]. In Ref. [23], the same analysis was repeated taking into account non-leading terms in the HQE. Finally, in [9, 24] also the full radiative corrections to $b \rightarrow cud$ were calculated and those parts of the corrections to $b \rightarrow ccs$ available from [7] were taken into account. Also the effect of scheme-dependence was estimated by changing the definition of the quark mass. For the same set of input parameters, the results are given in Table 1. The table shows that the introduction of nonperturbative correction terms by Bigi et al. reduces $B(B \rightarrow X e \nu)$ by 0.3% with respect to the free quark decay model, and that the account for the c quark mass in the radiative corrections yields an additional -0.9%, which again shows the importance of

perturbative corrections to the HQE. The scheme-dependence, however, is still tremendous and amounts to an uncertainty in $B(B \rightarrow X e \nu)$ of more than 1%. Whereas Refs. [22, 23] concluded that $B(B \rightarrow X e \nu) > 12.5\%$, Refs. [9, 24] find

$$B(B \rightarrow X e \nu) = (11.0 \pm 1.8 \pm 1.0)\%, \quad (8)$$

where the first error combines uncertainties in $\alpha_s(m_Z)$, m_b ($4.5 \text{ GeV} < m_b < 5.1 \text{ GeV}$), the hadronic corrections, the renormalization scale and the uncertainty in the radiative corrections to $b \rightarrow ccs$. The second error is a “guestimate” of the theoretical error due to scheme-dependence. The combined effect of complete radiative corrections, new results on $\alpha_s(m_Z)$ [25] and the consideration of different definitions of the quark mass thus lowers the theoretical branching ratio, which now agrees with the experimental one within the errors and seriously restricts any possible “new physics” in nonleptonic B decays. A more detailed analysis is in preparation [24].

References

- [1] I. Bigi, N. Uraltsev and A. Vainshtein, Phys. Lett. **B293** (1992) 430; *Erratum ibid.* **B297** (1993) 477;
I. Bigi *et al.*, Phys. Rev. Lett. **71** (1993) 496;
A. Manohar and M.B. Wise Phys. Rev. **D49** (1994) 1310;
B. Blok *et al.*, Phys. Rev. **D49** (1994) 3356;
T. Mannel, Nucl. Phys. **B413** (1994) 396.
- [2] M. Neubert, Phys. Rev. **D49** (1994) 3392;
T. Mannel and M. Neubert, Phys. Rev. **D50** (1994) 2037;
I. Bigi *et al.*, Int. J. Mod. Phys. **A9** (1994) 2467.
- [3] I. Bigi *et al.*, Minneapolis preprint: TPI-MINN-94/12-T (hep-ph/9405410).
- [4] I. Bigi *et al.*, Minneapolis preprint: TPI-MINN-94/25-T (hep-ph/9407296).
- [5] N. Cabibbo and L. Maiani, Phys. Lett. **B79** (1978) 109.
- [6] G. Altarelli *et al.*, Nucl. Phys. **B187** (1981) 461;
G. Buchalla, Nucl. Phys. **B391** (1993) 501.
- [7] Q. Hokim and X.Y. Pham, Phys. Lett. **B122** (1983) 297; Ann. Phys. **155** (1984) 202.
- [8] Y. Nir, Phys. Lett. **B221** (1989) 184.
- [9] E. Bagan *et al.*, TU München preprint: TUM-T31-67/94/R (hep-ph/9408306).
- [10] I. Bigi *et al.*, Phys. Rev. **D50** (1994) 2234;
M. Beneke and V.M. Braun, MPI München preprint: MPI-PhT/94-9 (hep-ph/9402364) (to appear in Nucl. Phys. **B**).

- [11] M. Beneke, V.M. Braun and V.I. Zakharov, MPI München preprint: MPI-PhT/94-18 (hep-ph/9405304).
- [12] P. Ball and V.M. Braun, Phys. Rev. D **49** (1994) 2472.
- [13] M. Neubert, Phys. Lett. B **322** (1994) 419.
- [14] M. Neubert, Talk given at QCD 94, Montpellier, France, 7-13 July 1994.
- [15] M. Shifman, N. Uraltsev and A. Vainshtein, Minneapolis preprint: TPI-MINN-94/13-T (hep-ph/9405207).
- [16] M. Luke and M. Savage, Phys. Lett. **B321** (1994) 88.
- [17] I. Bigi and N. Uraltsev, Z. Phys. **C62** (1994) 623.
- [18] Z. Ligeti and Y. Nir, Phys. Rev. **D49** (1994) 4331.
- [19] P. Ball and U. Nierste, TU München preprint: TUM-T31-56/94/R (hep-ph/9403407) (to appear in Phys. Rev. **D**).
- [20] Particle Data Group: M. Aguilar-Benitez *et al.*, Phys. Rev. **D50** (1994) 1173.
- [21] CLEO Collaboration: M.Athanas *et al.*, Cornell preprint: CLNS 94-1286 (hep-ex/9406004).
- [22] G. Altarelli and S. Petrarca, Phys. Lett. **B261** (1991) 303.
- [23] I. Bigi *et al.*, Phys. Lett. **B323** (1994) 408.
- [24] E. Bagan *et al.*, TU München preprint: TUM-T31-68/94 (in preparation).
- [25] S. Bethke, Talk given at QCD 94, Montpellier, France, 7-13 July 1994.